



Decision Support

Right-left asymmetry of the eigenvector method: A simulation study

László Csató^{a,b}^a HUN-REN Institute for Computer Science and Control (HUN-REN SZTAKI) Laboratory on Engineering and Management Intelligence Research Group of Operations Research and Decision Systems, Budapest, Hungary^b Corvinus University of Budapest (BCE) Institute of Operations and Decision Sciences Department of Operations Research and Actuarial Sciences, Budapest Hungary

ARTICLE INFO

Keywords:

Decision analysis
Analytic hierarchy process (AHP)
Eigenvector method
Right-left asymmetry
Simulation

ABSTRACT

The eigenvalue method, suggested by the developer of the extensively used Analytic Hierarchy Process methodology, exhibits right-left asymmetry: the priorities derived from the right eigenvector do not necessarily coincide with the priorities derived from the reciprocal left eigenvector. This paper offers a comprehensive numerical experiment to compare the two eigenvector-based weighting procedures and their reasonable alternative of the row geometric mean with respect to four measures. The underlying pairwise comparison matrices are constructed randomly with different dimensions and levels of inconsistency. The disagreement between the two eigenvectors turns out to be not always a monotonic function of these important characteristics of the matrix. The ranking contradictions can affect alternatives with relatively distant priorities. The row geometric mean is found to be almost at the midpoint between the right and inverse left eigenvectors, making it a straightforward compromise between them.

1. Introduction

“A suggestion for the prioritization of alternatives using the Perron-Frobenius right eigenvector of a pairwise comparison matrix has recently been made by T. Saaty. We note that use of the left eigenvectors is equally justified (as long as order is reversed).”¹

The Analytic Hierarchy Process (AHP) is one of the most popular decision-making techniques since it has been introduced by Saaty (1977, 1980). It has a number of successful applications (Bhushan & Rai, 2007; Forman & Gass, 2001; Vaidya & Kumar, 2006; Vargas, 1990) and, simultaneously, several flaws identified in the literature (Csató, 2017; Csató & Petróczy, 2021; Genest, Lapointe, & Drury, 1993; Munier & Hontoria, 2021; Petróczy & Csató, 2021).

The current paper deals with an issue in priority derivation from a given pairwise comparison matrix. Saaty has suggested using the right eigenvector for this purpose but there are many other methods (Choo & Wedley, 2004). In particular, Johnson et al. (1979) argue for the componentwise reciprocal of the left eigenvector as written in the motto above. Another strong competitor is the logarithmic least squares or row geometric mean (Crawford & Williams, 1985), mainly due to its strong axiomatic foundations (Barzilai, 1997; Barzilai, Cook, & Golany, 1987; Bozóki & Tsyganok, 2019; Csató, 2018; 2019; Fichtner, 1984; 1986; Lundy, Siraj, & Greco, 2017).

In the following, these solutions will be compared using a Monte Carlo simulation approach, that is, the priority vectors of the three weighting procedures are evaluated on the basis of a large set of random pairwise comparison matrices. Even though a similar exercise has been attempted at least twice in the previous literature (Bozóki & Rapcsák, 2008; Ishizaka & Lusti, 2006), independently of each other, we make important contributions compared to both preliminary studies:

- Ishizaka & Lusti (2006) limit the investigation to at most seven alternatives and matrices with an acceptable level of inconsistency. The simulation has some drawbacks because (1) the number of matrices for a given order is 500, which might be insufficient to derive robust results; and (2) the matrix entries can only be integers. Last but not least, the authors focus exclusively on the number of ranking contradictions and do not consider the differences in the priorities derived from the three methods.
- Bozóki & Rapcsák (2008) examine the case of five alternatives. All matrix entries are chosen randomly, hence, the number of matrices with an acceptable level of inconsistency remains rather low. The authors only consider the frequency of rank reversals.

Here, 3 million pairwise comparison matrices are generated for any given number of alternatives between four and nine such that each

E-mail address: laszlo.csato@sztaki.hu.

¹ Source: Johnson, Beine, & Wang (1979, p. 61).

interval of Saaty’s inconsistency ratio (Saaty, 1977) with a length of 0.005 contains at least one thousand instances. The right and inverse left eigenvectors, as well as the row geometric mean, are evaluated according to four measures: the Euclidean and Chebyshev distances, as well as the Kendall rank correlation coefficients of the normalised weight vectors, and the maximal ratios of the priorities corresponding to one alternative. Among them, only Kendall rank correlation depends on the ranking of the alternatives.

Our comprehensive analysis yields several interesting results, some of them at least partially contradicting previous findings:

- The row geometric mean is revealed to be an excellent compromise between the right and inverse left eigenvectors as it is almost at the midpoint between them, especially at a low level of inconsistency;
- The differences between the three priority deriving methods do not always increase with the level of inconsistency if the latter is relatively high (we refine the conclusion of Ishizaka & Lusti (2006, p. 398));
- The differences between the three priority deriving methods do not always increase with the number of alternatives (we refine the conclusion of Ishizaka & Lusti (2006, p. 398));
- Three examples illustrate that (1) rank reversal between the right and inverse left eigenvectors may emerge for a slightly perturbed consistent matrix; (2) the right and inverse left eigenvectors can lead to a fully reversed order of the alternatives; (3) rank reversal between the right and inverse left eigenvectors might occur even if the priorities of two alternatives are distant (this denies the conclusion of Ishizaka & Lusti (2006, p. 398)).

The main reason for the different conclusions compared to Ishizaka & Lusti (2006) resides in the extension of our analysis to (a) a wider interval of inconsistency (inconsistency ratio $CR < 0.5$ rather than $CR < 0.1$); (b) a higher number of alternatives (up to nine instead of seven); (c) a broader set of comparison metrics. The expansion of the range of inconsistency can be justified since $CR < 0.1$ is an inflexible criterion and is too restrictive when the size of the matrix increases (Alonso & Lamata, 2006). Furthermore, the results are based on a much higher number of random matrices.

The remainder of the study is organised as follows. The mathematical background is presented in Section 2. Section 3 reviews the related literature, and Section 4 outlines the simulation experiment. Section 5 contains the main results. The paper is finished with a concise discussion in Section 6.

2. Theoretical background

An $n \times n$ matrix $A = [a_{ij}]$ is called a pairwise comparison matrix if it is positive ($a_{ij} > 0$ for all i, j) and reciprocal ($a_{ji} = 1/a_{ij}$ for all i, j). Its entry a_{ij} quantifies how many times alternative i is better/more important compared to alternative j . An important property of a pairwise comparison matrix is *consistency*: it is called consistent if $a_{ik} = a_{ij}a_{jk}$ for all i, j, k ; otherwise, it is called inconsistent.

Pairwise comparison matrices are mainly used to derive priorities for the alternatives. In the case of a consistent matrix, this is almost trivial since the matrix is generated by an appropriate weight vector, that is, there exists a vector $w = [w_i]$ such that $a_{ij} = w_i/w_j$ for all i, j . For inconsistent matrices, several weighting techniques have been proposed in the literature (Choo & Wedley, 2004). Probably the two most popular procedures are the logarithmic least squares (LLSM) or row geometric mean (Crawford & Williams, 1985; De Graan, 1980; de Jong, 1984; Rabinowitz, 1976; Williams & Crawford, 1980) and the eigenvector (Saaty, 1977) methods.

The logarithmic least squares method minimises the aggregated distances of the approximations in a logarithmic sense:

$$\sum_{i=1}^n \sum_{j=1}^n \left[\log(a_{ij}) - \log\left(\frac{w_i}{w_j}\right) \right]^2 \rightarrow \min.$$

The solution is provided by the geometric means of row elements, namely:

$$\frac{w_i}{w_j} = \frac{\sqrt[n]{\prod_{k=1}^n a_{ik}}}{\sqrt[n]{\prod_{k=1}^n a_{jk}}}$$

The corresponding weight vector is denoted by w^{RGM} .

The eigenvector method is based on the right eigenvector associated with the dominant eigenvalue λ_{\max} of the pairwise comparison matrix:

$$A w^R = \lambda_{\max} w^R.$$

However, the matrix has also a left eigenvector associated with the same dominant eigenvalue λ_{\max} :

$$w^L A = \lambda_{\max} w^L.$$

If the pairwise comparison matrix is consistent, the componentwise inverse of the left eigenvector is the right eigenvector: $w_i^L = 1/w_i^R$. Therefore, it is reasonable to use the componentwise inverse of the left eigenvector, denoted by w^{-L} in the following, to derive the priorities (Johnson et al., 1979).

Finally, the (geometric) mean of the right and inverse left eigenvectors $w^{RL} = [w_i^{RL}]$ can be defined as

$$w_i^{RL} = w_i^R w_i^{-L}$$

for all $1 \leq i \leq n$.

Several inconsistency measures have been suggested in the literature (Brunelli, 2018). In this paper, we use the first index proposed by Saaty (1977):

$$CI = \frac{\lambda_{\max} - n}{n - 1}.$$

The value of CI is compared to the average CI of a high number of randomly generated pairwise comparison matrices, which is denoted by RI , in order to get the inconsistency ratio $CR = CI/RI$. According to Saaty, a pairwise comparison matrix can be accepted if CR does not exceed 0.1. A statistical interpretation of the 10% rule is provided by Vargas (1982).

3. The significance of the problem

A conceptual weakness of the eigenvector method is the issue of asymmetry (Bozóki & Rapcsák, 2008). The entry a_{ij} of a pairwise comparison matrix A gives the numerical answer to the question “How much does alternative i dominate alternative j ?”. However, one can equivalently ask the reciprocal question of “How much does alternative j dominate alternative i ?” to arrive at matrix A^T . The latter approach produces the right eigenvector of A^T , which is the elementwise reciprocal of the right eigenvector of A .

A crucial motivation behind the eigenvector method comes from the property of consistent pairwise comparison matrices that $a_{ij} = w_i/w_j$ and $w_i = a_{ij}w_j$ for all i, j . This system of linear equations leads almost directly to the matrix equation $\lambda w = Aw$. However, $a_{ij} = w_i/w_j$ can be equivalently written as $w_j = a_{ji}w_i$ for all i, j , which implies the matrix equation $A^T w = \lambda w$. Obviously, the right eigenvector of A^T is the left eigenvector of A . However, the meaning of the entries in A^T is the opposite compared to the entries of A . Consequently, “we might just as well use a left eigenvector to prioritize in the general case as long as order reversal is allowed for” (Johnson et al., 1979, p. 62).

It might also happen that the decision-maker misinterprets the task,




Fig. 1. The preferences of two decision-makers in Example 1.

and provides all pairwise comparisons in a reversed order (Dodd, Donegan, & McMaster, 1995). This reversal results in the emergence of the left rather than the right eigenvector. Hence, “if these two vectors are not componentwise mutual inverses, it is impossible to say which is ‘correct’ (Dodd et al., 1995, p. 88).”

In order to highlight further why the right-left asymmetry can be important, a simple group decision-making problem with two sets of preferences is considered.

Example 1. Fig. 1 presents the numerical preferences of two decision-makers (DMs) for four alternatives. For instance, alternative S is judged to be twice more important as alternative T by DM1, but S is only half more important compared to T according to DM2. Indeed, the preferences of DM1 and DM2 are exactly the opposite. Therefore, it is natural to assign the same priorities to all alternatives S, T, U, V on the basis of the aggregated preferences of the two DMs.

Take the associated pairwise comparison matrices **A** and **B** of DM1 and DM2, respectively:

$$A = \begin{bmatrix} 1 & 1 & 1 & 9 \\ 1 & 1 & 2 & 5 \\ 1 & 1/2 & 1 & 9 \\ 1/9 & 1/5 & 1/9 & 1 \end{bmatrix} \quad B = \begin{bmatrix} 1 & 1 & 1 & 1/9 \\ 1 & 1 & 1/2 & 1/5 \\ 1 & 2 & 1 & 1/9 \\ 9 & 5 & 9 & 1 \end{bmatrix}.$$

The corresponding right eigenvectors are as follows (the sum of priorities are normalised to 100):

$$w^R(A) = [32.42 \quad 35.02 \quad 28.21 \quad 4.35];$$

$$w^R(B) = [8.86 \quad 9.05 \quad 11.04 \quad 71.05].$$

In group decision-making, there are essentially two ways to derive aggregated priorities for the alternatives: (1) aggregating the individual pairwise comparison matrices, and using a weighting method for the aggregated matrix (Aczél & Saaty, 1983); or (2) using a weighting method for the individual pairwise comparison matrices, and aggregating these priorities (Basak & Saaty, 1993). Aczél & Saaty (1983) provide an axiomatic approach to show that individual matrices should be aggregated by the geometric mean when the aggregation of **A** and **B** = **A**^T results in a consistent matrix in which all entries are one. Then each alternative should have the same priority.

On the other hand, any reasonable aggregation procedure of **w**^R(**A**) and **w**^R(**B**) results in a higher weight for alternative T than for alternative S since the second entry of both individual weight vectors is higher than the first entry.

Note that in this case, the disturbingly different implications of techniques (1) and (2) are caused exclusively by the difference between the right and inverse left eigenvectors.

According to Example 1, a potential cause of the well-known difference between aggregation procedures (1) and (2) for the eigenvector method can be the right-left asymmetry. Thus, using the right

eigenvector can be strongly debated in group decision-making if the right and inverse left eigenvectors differ. Based on these arguments, we agree with Johnson et al. (1979, p. 62) that “there is no reason to believe that utilization of a right eigenvector yields a ‘better’ scheme than the left.”

4. Literature review

While the components in the normalised left eigenvector of a pairwise comparison matrix with three alternatives are the reciprocals of the components of the normalised right eigenvector, this is not necessarily the case when the number of alternatives is at least four. This problem has been identified first by Johnson et al. (1979), who presented the following example:

$$A = \begin{bmatrix} 1 & 3 & 1/3 & 1/2 \\ 1/3 & 1 & 1/6 & 2 \\ 3 & 6 & 1 & 1 \\ 2 & 1/2 & 1 & 1 \end{bmatrix}.$$

Here, the right eigenvector—if the sum of priorities equals 100—is:

$$w^R(A) = [18.44 \quad 15.19 \quad 43.64 \quad 22.73].$$

Consequently, the fourth alternative is ranked above the first.

The left eigenvector is

$$w^L(A) = [24.82 \quad 38.78 \quad 10.49 \quad 25.91],$$

hence, the elementwise reciprocal left eigenvector is

$$w^{-L}(A) = [20.14 \quad 12.89 \quad 47.67 \quad 19.29],$$

implying that the first alternative is preferred to the fourth.

The consistency ratio of matrix **A** is $CR(A) \approx 0.331$, which cannot be accepted according to Saaty’s 10% rule.

Johnson et al. (1979) have also generated 364 random matrices of order six with $CI < 1$ and found 164 ranking interchanges between the right and the componentwise reciprocal left eigenvectors. In addition, the authors note that the disagreement can occur for arbitrarily small positive values of CR due to continuity.

For $n \geq 4$, the reciprocal property between the left and right eigenvector components holds not only if the matrix is consistent (DeTurck, 1987). In the case of the pairwise comparison matrix

$$B = \begin{bmatrix} 1 & 8/5 & 1/4 & 4 \\ 5/8 & 1 & 5/8 & 10 \\ 4 & 8/5 & 1 & 4 \\ 1/4 & 1/10 & 1/4 & 1 \end{bmatrix},$$

the right eigenvector is

$$w^R(B) = [2/9 \quad 5/18 \quad 4/9 \quad 1/18],$$

and the left eigenvector is

$$w^L(B) = [1/4 \quad 1/5 \quad 1/8 \quad 1],$$

thus, $w^{-L}(A) = w^R(A)$. However, $CR(B)$ is higher than 0.1.

An acceptable inconsistency ratio does not guarantee that the right and inverse left eigenvectors imply the same ranking of the alternatives (Dodd et al., 1995). Let

$$C = \begin{bmatrix} 1 & 1 & 3 & 9 & 9 \\ 1 & 1 & 5 & 8 & 5 \\ 1/3 & 1/5 & 1 & 9 & 5 \\ 1/9 & 1/8 & 1/9 & 1 & 1 \\ 1/9 & 1/5 & 1/5 & 1 & 1 \end{bmatrix},$$

where $CR(C) \approx 0.082$. The priorities from the right eigenvector are

$$w^R(C) = [36.5652 \quad 38.9564 \quad 16.7155 \quad 3.4693 \quad 4.2936],$$

and the priorities from the componentwise reciprocal of the left eigenvector are

$$w^{-L}(C) = [40.6431 \quad 36.4208 \quad 15.0669 \quad 3.4391 \quad 4.4302].$$

Therefore, the first alternative is the best by the left eigenvector but the second should be chosen by the right eigenvector.

Ishizaka & Lusti (2006) compare all the three weighting techniques—logarithmic least squares method, right, and inverse left eigenvectors—defined in Section 2 for five intervals of the inconsistency ratio (0–0.02, 0.02–0.04, 0.04–0.06, 0.06–0.08, 0.08–0.1) and three to seven alternatives. The underlying matrices are generated randomly. Ranking contradictions are found to increase linearly with the level of inconsistency and the dimension of the matrix. However, the number of matrices in each case (100) is rather low, the matrix entries are integers, and the authors focus only on the ranking of the alternatives.

Bozóki & Rapsák (2008, Section 6) examine 100 million randomly generated pairwise comparison matrices of order five to estimate the frequency of rank reversal between the weights computed from the right and inverse left eigenvectors. This measure is found to increase along with the inconsistency ratio CR . However, the entries of the pairwise comparison matrix are chosen independently of each other, thus, the number of matrices for small values of CR is relatively small. Furthermore, it remains to be seen whether the same result holds if the number of alternatives varies.

Tomashevskii (2015) argues that the derived weights cannot be used to rank the alternatives without taking their errors into account, and the rank reversal between the right and the inverse left eigenvectors occurs only because the errors are high. Consequently, the means do not contain any information on the ranking of the corresponding alternatives.

5. The design of numerical experiments

Our matrix generation algorithm is based on constructing a consistent pairwise comparison matrix and perturbing its elements. The technique of Szádóczki, Bozóki, Juhász, Kadenko, & Tsyganok (2023) is followed as it improves the method of Szádóczki, Bozóki, & Tekile (2022). This consists of the following steps:

1. Choosing n uniformly distributed random number w_i from the interval $[1; 9]$. Computing a consistent pairwise comparison matrix $A = [a_{ij}]$, where $a_{ij} = w_i/w_j$ for all i, j .

2. For all $i \neq j$, either a_{ij} or a_{ji} is perturbed depending on which entry is higher.

If $a_{ij} \geq 1$, then the perturbed entry \hat{a}_{ij} is

$$\hat{a}_{ij} = \begin{cases} a_{ij} + \varepsilon_{ij} & \text{if } a_{ij} + \varepsilon_{ij} \geq 1 \\ 1/[1 - \varepsilon_{ij} - (a_{ij} - 1)] & \text{otherwise,} \end{cases} \quad (1)$$

where ε_{ij} is a uniformly distributed random number from the interval $[-\Delta; \Delta]$.

If $a_{ij} < 1$, then $a_{ji} > 1$, and \hat{a}_{ji} is computed analogously to (1).

3. The reciprocity of the matrix is kept by adjusting the pair of the perturbed entry.

This process ensures that the perturbed elements are uniformly distributed around the original a_{ij} on the scale where the distance between $1/b$ and $1/c$ equals the distance between b and c , see Szádóczki et al. (2023, Fig. 1).

We consider three different values of parameter Δ (1, 2, 3) and six dimensions ($4 \leq n \leq 9$) for which 1 million matrices are generated, respectively (altogether 18 million). For any matrix, the inconsistency ratio and the weight vectors according to the three techniques presented in Section 2 are calculated.

The distribution of the pairwise comparison matrices with respect to the level of inconsistency is shown in Fig. 2. If $\Delta = 1$, almost all matrices have an inconsistency ratio below 0.1, except for $n = 4$. As the dimension of the matrix increases, the curves for a given Δ are more peaked and less asymmetric, meaning that inconsistency is more strongly determined by the maximal perturbation. This crucial observation is not reported in Szádóczki et al. (2023) since the authors provide only the average value of CR . The reader is directed to Szádóczki et al. (2023) for other characteristics of the simulated matrices.

In order to compare two priority vectors u and v normalised by $\sum_{i=1}^n u_i = 1$ and $\sum_{i=1}^n v_i = 1$, respectively, four metrics are considered:

- Euclidean distance:

$$d_{\text{euc}}(u, v) = \sqrt{\sum_{i=1}^n (u_i - v_i)^2}; \quad (2)$$

- Chebyshev distance:

$$d_{\text{cheb}}(u, v) = \max\{|u_i - v_i| : 1 \leq i \leq n\}; \quad (3)$$

- Maximal ratio:

$$\omega(u, v) = \max\left\{\max\left\{\frac{u_i}{v_i}, \frac{v_i}{u_i}\right\} : 1 \leq i \leq n\right\}; \quad (4)$$

- Kendall tau:

$$\tau(u, v) = \frac{\#c(u, v) - \#d(u, v)}{n(n-1)/2}; \quad (5)$$

where $\#c$ ($\#d$) denotes the number of (dis)concordant pairs $1 \leq i, j \leq n$ when $u_i > u_j$ and $v_i > v_j$ ($v_i < v_j$).

The Euclidean distance (2) is the length of a line segment between the two vectors. The Chebyshev distance (3) depends only on the greatest difference along any coordinate.

The maximal ratio (4) is inspired by the Chebyshev distance but focuses on ratios instead of differences since the Chebyshev distance does not reflect high relative deviations in the weights of lower-ranked alternatives. For instance, let $u = [0.5; 0.4; 0.1]$, $v = [0.5; 0.3; 0.2]$, and




Fig. 2. The distribution of randomly generated matrices according to their inconsistency ratio CR (sample size: 10^6 in each case; resolution of the intervals: 0.005).

$w = [0.6; 0.3; 0.1]$. Then $d_{cheb}(u, v) = d_{cheb}(u, w) = 0.1$ but $\omega(u, v) = 2 > 4/3 = \omega(u, w)$. Clearly, the Chebyshev distance does not take into account whether the switching in priorities happens between the top two (u and w) or the bottom two (u and v) alternatives. However, the latter is more serious in terms of relative changes, which might be important if the priorities are used, for example, to allocate some resources proportionally.

Finally, Kendall tau (5) is a standard rank correlation coefficient (Kendall, 1938). Its range is $[-1; 1]$ with -1 showing two opposite rankings and $+1$ indicating two identical rankings. In contrast to the other three measures, here a higher value is associated with a stronger agreement between the two weight vectors.

6. Results

For each of the four metrics (2)–(5) and each interval of the inconsistency ratio CR (resolution: 0.005), we compute

- the average value of the metric between the right eigenvector w^R and the inverse left eigenvector w^{-L} for all pairwise comparison matrices at the given level of inconsistency;
- the average value of the metric between the right eigenvector w^R and the (geometric) mean of the right and inverse left eigenvectors w^{RL} for all pairwise comparison matrices at the given level of inconsistency;
- the average value of the metric between the right eigenvector w^R and the row geometric mean weight vector w^{RGM} for all pairwise comparison matrices at the given level of inconsistency;
- the probability that the row geometric mean weight vector w^{RGM} is not farther from the right eigenvector w^R than the inverse left eigenvector w^{-L} , based on all pairwise comparison matrices at the given level of inconsistency.

In order to ensure the robustness of the results, only the inconsistency intervals with at least one thousand matrices are shown; this accounts for the smaller range in the case of higher dimensions (cf. Fig. 2).

Fig. 3 uses the Euclidean, while Fig. 4 follows the Chebyshev approach to quantify the distances of the weights. In the case of average




Fig. 3. The Euclidean distances of weight vectors as a function of the inconsistency ratio CR (resolution of the intervals: 0.005).

distances, the shapes of the lines are almost indistinguishable for a given n . For small levels of inconsistency, the distances between the right and inverse left eigenvectors increase almost linearly as a function of inconsistency, which coincides with the finding of Ishizaka & Lusti (2006). However, the growth slackens and the maximum distance is reached for a given value of CR somewhere between 0.22 (if $n = 9$) and 0.4 (if $n = 5$). The only exception is $n = 4$, when a higher inconsistency is associated with a higher distance between the two eigenvectors.

On the other hand, the distance between the (right) eigenvector and the row geometric mean increases monotonically, albeit the curve is almost flat at higher levels of inconsistency. If CR does not exceed 10%, that is, inconsistency can be accepted according to the criterion of Saaty, then the logarithmic least squares method is essentially at the midpoint between the right and inverse left eigenvectors, and is close to the geometric mean of the two eigenvectors. This seems to be a quite powerful argument for using the row geometric mean method to derive the weights.

Since the higher average distance between the two eigenvectors \mathbf{w}^R and \mathbf{w}^{-L} compared to the average distance between \mathbf{w}^R and the row

geometric mean \mathbf{w}^{RGM} may be misleading, the probability that \mathbf{w}^R is closer to \mathbf{w}^{RGM} than to \mathbf{w}^{-L} has also been calculated. If $CR < 0.1$, this is almost guaranteed for the Chebyshev distance, while the likelihood is still above 98% for the Euclidean distance. The probability exceeds 80% even at higher levels of inconsistency, and it is increasing with the number of alternatives at a given value of CR , except for $4 \leq n \leq 5$ and the Chebyshev distance.

Fig. 5, which focuses on the mean maximal ratios of the priorities, reinforces the findings from Figs. 3 and 4. Consequently, the cardinal difference between the right and inverse left eigenvectors, as well as between the (right) eigenvector and the row geometric mean is robust and almost independent of the measure chosen.

In contrast to the previous three metrics, Kendall tau considers only the rankings implied by the weights. Since a higher value shows a stronger similarity, Fig. 6 plots the difference of the average Kendall tau from its theoretical maximum of one. The number of ranking contradictions generally increases along with the level of inconsistency, but two breaks appear in the lines, especially for higher n , which are probably caused by the three different values for parameter Δ . Ishizaka &




Fig. 4. The Chebyshev distances of weight vectors as a function of the inconsistency ratio CR (resolution of the intervals: 0.005).

Lusti (2006, Fig. 3) present a similar result based on a much smaller number of randomly generated matrices.

However, according to Ishizaka & Lusti (2006, Fig. 4), the ranking contradiction phenomenon increases linearly with the dimension of the matrix because the possibility of a reversal rises if there are more alternatives. Fig. 6 does not fully support this conclusion: the value of mean Kendall tau is about the same for a given inconsistency interval if the number of alternatives is at least seven (e.g. about 0.9 if CR is approximately 0.1). Again, the difference between the (right) eigenvector and the row geometric mean is only slightly larger than the difference between the right eigenvector and the (geometric mean) of the two eigenvectors. In addition, the ranking implied by the row geometric mean is not farther from the ranking implied by the right eigenvector than the ranking implied by the left eigenvector with a probability of at least 95% (except for $n = 4$, where this remains true only if $CR < 0.2$). Hence, the row geometric mean seems to be a reasonable compromise between the two eigenvectors even from an ordinal point of view.

Besides the results based on a high number of random pairwise comparison matrices, our simulations have provided some interesting

examples that are worth further consideration. First, according to Johnson et al. (1979, p. 63), the disagreement between the right and inverse left eigenvectors can occur at arbitrarily small positive values of inconsistency. For the pairwise comparison matrix

$$\mathbf{M}^{(1)} = \begin{bmatrix} 1 & 0.4759 & 0.9832 & 0.4025 \\ 2.1011 & 1 & 1.9975 & 0.7374 \\ 1.0171 & 0.5006 & 1 & 0.3704 \\ 2.4842 & 1.3560 & 2.6998 & 1 \end{bmatrix},$$

the weights from the right eigenvector are

$$\mathbf{w}^R(\mathbf{M}^{(1)}) = [15.042 \quad 30.274 \quad 15.037 \quad 39.647],$$

and the weights from the componentwise reciprocal of the left eigenvector are

$$\mathbf{w}^{-L}(\mathbf{M}^{(1)}) = [15.036 \quad 30.281 \quad 15.049 \quad 39.635].$$




Fig. 5. The maximal ratios of weight vectors as a function of the inconsistency ratio CR (resolution of the intervals: 0.005).

Rank reversal arises between the first and the third alternatives in the last two positions, although $CR(\mathbf{M}^{(1)}) \approx 0.0007$. Therefore, a class of pairwise comparison matrices with minimal inconsistency can be sought such that the right and inverse left eigenvectors lead to a different ranking of the alternatives, similar to the issue of Pareto inefficiency (Bozóki, 2014).

Second, the left and right eigenvector components might imply an opposite order of the alternatives. In the case of the pairwise comparison matrix

$$\mathbf{M}^{(2)} = \begin{bmatrix} 1 & 1.624 & 0.574 & 1.072 & 1.054 \\ 0.616 & 1 & 1.132 & 1.089 & 1.269 \\ 1.743 & 0.884 & 1 & 1.515 & 0.467 \\ 0.933 & 0.919 & 0.660 & 1 & 1.694 \\ 0.949 & 0.788 & 2.140 & 0.590 & 1 \end{bmatrix},$$

the priorities from the right eigenvector are

$$\mathbf{w}^R(\mathbf{M}^{(2)}) = [19.75 \quad 19.16 \quad 20.85 \quad 19.53 \quad 20.71],$$

and the priorities from the componentwise reciprocal of the left eigenvector are

$$\mathbf{w}^{-L}(\mathbf{M}^{(2)}) = [20.25 \quad 20.55 \quad 19.31 \quad 20.27 \quad 19.62].$$

Hence, the ranking of the alternatives is $3 \succ 5 \succ 1 \succ 4 \succ 2$ in the former case, which is reversed to $2 \succ 4 \succ 1 \succ 5 \succ 3$ in the latter case. The inconsistency ratio is $CR(\mathbf{M}^{(2)}) \approx 0.078$.

Third, it has been thought that “Only very close priorities suffer from ranking contradictions” (Ishizaka & Lusti, 2006, p. 398). Even though the exact meaning of very close remains obscure, the following example probably disproves this statement:




Fig. 6. The Kendall rank correlation coefficients of weight vectors as a function of the inconsistency ratio CR (resolution of the intervals: 0.005).

$$\mathbf{M}^{(3)} = \begin{bmatrix} 1 & 0.371 & 2.013 & 5.389 & 0.243 \\ 2.698 & 1 & 4.596 & 7.527 & 0.736 \\ 0.497 & 0.218 & 1 & 2.321 & 0.167 \\ 0.186 & 0.133 & 0.431 & 1 & 0.385 \\ 4.120 & 1.359 & 5.973 & 2.598 & 1 \end{bmatrix},$$

where the weights from the right eigenvector are

$$\mathbf{w}^R(\mathbf{M}^{(2)}) = [15.26 \quad 33.23 \quad 7.74 \quad 5.68 \quad 38.08],$$

and the weights from the componentwise reciprocal of the left eigenvector are

$$\mathbf{w}^{-L}(\mathbf{M}^{(2)}) = [15.29 \quad 37.84 \quad 8.55 \quad 4.93 \quad 33.39].$$

Consequently, the best alternative is the fifth according to the right and the second according to the inverse left eigenvector. The absolute value of the difference between the weights of the second and the fifth alter-

natives is 4.85 and 4.44, respectively, if the sum of weights is normalised to 100. The inconsistency ratio is $CR(\mathbf{M}^{(3)}) \approx 0.0993$.

7. Conclusions

The paper has addressed one of the most serious shortcomings of the eigenvector method, a widely used priority deriving technique for pairwise comparison matrices. In particular, we have compared the weights implied by the right and inverse left eigenvectors, as well as by the row geometric mean for a large set of randomly generated matrices. The two eigenvectors turned out to lead to different priorities and rankings relatively often, and their disagreement is not necessarily a monotonic function of the level of inconsistency and the number of alternatives. Although the problem is less threatening if inconsistency remains below the acceptable threshold, the left and right eigenvector components may imply an opposite priority order or a rank reversal between alternatives with distant weights even if the value of the inconsistency ratio does not exceed 10%.

There are at least two important conclusions to be drawn from these

results. Our findings uncover that the row geometric mean is almost at the midpoint between the principal right and inverse left eigenvectors, hence, it is a reasonable compromise between them. Therefore, a novel argument is provided for following this method, which is also easy to calculate and satisfies many attractive theoretical properties. Furthermore, we have reinforced an important message of Johnson et al. (1979): assuming the existence of a single ranking or priority vector is usually too demanding. Rather, a range of possible orderings and weights can be allowed (instead of a single ranking and exact priorities) such that the range is wider for a higher level of inconsistency and uncertainty in the input data.

Finally, this research is far from finished and can be continued in several directions. First, the disagreement between the two eigenvectors is worth analysing on particular classes of pairwise comparison matrices. For instance, following the work of DeTurck (1987), it would be interesting to characterise the set of matrices for which the reciprocal property between the left and right eigenvector components hold. Second, a new weighting method can be introduced by aggregating the two eigenvectors appropriately. Third, other deficiencies of the eigenvector method, such as Pareto inefficiency (Blanquero, Carrizosa, & Conde, 2006; Bozóki & Fülöp, 2018), might be studied in a similar Monte Carlo experiment.

Acknowledgments

We are grateful to Sándor Bozóki and Zsombor Szádóczki for useful advice.

Three anonymous reviewers provided valuable comments and suggestions on earlier drafts.

References

- Aczél, J., & Saaty, T. L. (1983). Procedures for synthesizing ratio judgements. *Journal of Mathematical Psychology*, 27(1), 93–102.
- Alonso, J. A., & Lamata, M. T. (2006). Consistency in the analytic hierarchy process: A new approach. *International Journal of Uncertainty, Fuzziness and Knowledge-Based Systems*, 14(4), 445–459.
- Barzilai, J. (1997). Deriving weights from pairwise comparison matrices. *Journal of the Operational Research Society*, 48(12), 1226–1232.
- Barzilai, J., Cook, W. D., & Golany, B. (1987). Consistent weights for judgements matrices of the relative importance of alternatives. *Operations Research Letters*, 6(3), 131–134.
- Basak, I., & Saaty, T. (1993). Group decision making using the analytic hierarchy process. *Mathematical and Computer Modelling*, 17(4–5), 101–109.
- Bhushan, N., & Rai, K. (2007). *Strategic Decision Making: Applying the Analytic Hierarchy Process*. London: Springer Science & Business Media.
- Blanquero, R., Carrizosa, E., & Conde, E. (2006). Inferring efficient weights from pairwise comparison matrices. *Mathematical Methods of Operations Research*, 64(2), 271–284.
- Bozóki, S. (2014). Inefficient weights from pairwise comparison matrices with arbitrarily small inconsistency. *Optimization*, 63(12), 1893–1901.
- Bozóki, S., & Fülöp, J. (2018). Efficient weight vectors from pairwise comparison matrices. *European Journal of Operational Research*, 264(2), 419–427.
- Bozóki, S., & Rapcsák, T. (2008). On Saaty's and Koczkodaj's inconsistencies of pairwise comparison matrices. *Journal of Global Optimization*, 42(2), 157–175.
- Bozóki, S., & Tsyganok, V. (2019). The (logarithmic) least squares optimality of the arithmetic (geometric) mean of weight vectors calculated from all spanning trees for incomplete additive (multiplicative) pairwise comparison matrices. *International Journal of General Systems*, 48(4), 362–381.
- Brunelli, M. (2018). A survey of inconsistency indices for pairwise comparisons. *International Journal of General Systems*, 47(8), 751–771.
- Choo, E. U., & Wedley, W. C. (2004). A common framework for deriving preference values from pairwise comparison matrices. *Computers & Operations Research*, 31(6), 893–908.
- Crawford, G., & Williams, C. (1985). A note on the analysis of subjective judgment matrices. *Journal of Mathematical Psychology*, 29(4), 387–405.
- Csató, L. (2017). Eigenvector method and rank reversal in group decision making revisited. *Fundamenta Informaticae*, 156(2), 169–178.
- Csató, L. (2018). Characterization of the row geometric mean ranking with a group consensus axiom. *Group Decision and Negotiation*, 27(6), 1011–1027.
- Csató, L. (2019). A characterization of the logarithmic least squares method. *European Journal of Operational Research*, 276(1), 212–216.
- Csató, L., & Petróczy, D. G. (2021). On the monotonicity of the eigenvector method. *European Journal of Operational Research*, 292(1), 230–237.
- De Graan, J. G. (1980). Extensions of the multiple criteria analysis method of T. L. Saaty. *Report*. Voorburg: National Institute for Water Supply.
- de Jong, P. (1984). A statistical approach to Saaty's scaling method for priorities. *Journal of Mathematical Psychology*, 28(4), 467–478.
- DeTurck, D. M. (1987). The approach to consistency in the analytic hierarchy process. *Mathematical Modelling*, 9(3–5), 345–352.
- Dodd, F. J., Donegan, H. A., & McMaster, T. B. M. (1995). Inverse inconsistency in analytic hierarchies. *European Journal of Operational Research*, 80(1), 86–93.
- Fichtner, J. (1984). Some thoughts about the mathematics of the Analytic Hierarchy Process. *Technical Report*. Institut für Angewandte Systemforschung und Operations Research, Universität der Bundeswehr München.
- Fichtner, J. (1986). On deriving priority vectors from matrices of pairwise comparisons. *Socio-Economic Planning Sciences*, 20(6), 341–345.
- Forman, E. H., & Gass, S. I. (2001). The analytic hierarchy process—an exposition. *Operations Research*, 49(4), 469–486.
- Genest, C., Lapointe, F., & Drury, S. W. (1993). On a proposal of Jensen for the analysis of ordinal pairwise preferences using Saaty's eigenvector scaling method. *Journal of Mathematical Psychology*, 37(4), 575–610.
- Ishizaka, A., & Lusti, M. (2006). How to derive priorities in AHP: A comparative study. *Central European Journal of Operations Research*, 14(4), 387–400.
- Johnson, C. R., Beine, W. B., & Wang, T. J. (1979). Right-left asymmetry in an eigenvector ranking procedure. *Journal of Mathematical Psychology*, 19(1), 61–64.
- Kendall, M. G. (1938). A new measure of rank correlation. *Biometrika*, 30(1/2), 81–93.
- Lundy, M., Siraj, S., & Greco, S. (2017). The mathematical equivalence of the “spanning tree” and row geometric mean preference vectors and its implications for preference analysis. *European Journal of Operational Research*, 257(1), 197–208.
- Munier, N., & Hontoria, E. (2021). *Uses and Limitations of the AHP Method: A Non-Mathematical and Rational Analysis*. Cham, Switzerland: Springer.
- Petróczy, D. G., & Csató, L. (2021). Revenue allocation in Formula One: A pairwise comparison approach. *International Journal of General Systems*, 50(3), 243–261.
- Rabinowitz, G. (1976). Some comments on measuring world influence. *Conflict Management and Peace Science*, 2(1), 49–55.
- Saaty, T. L. (1977). A scaling method for priorities in hierarchical structures. *Journal of Mathematical Psychology*, 15(3), 234–281.
- Saaty, T. L. (1980). *The Analytic Hierarchy Process: Planning, Priority Setting, Resource Allocation*. New York: McGraw-Hill.
- Szádóczki, Z., Bozóki, S., Juhász, P., Kadenko, S. V., & Tsyganok, V. (2023). Incomplete pairwise comparison matrices based on graphs with average degree approximately 3. *Annals of Operations Research*, 326(2), 783–807.
- Szádóczki, Z., Bozóki, S., & Tekile, H. A. (2022). Filling in pattern designs for incomplete pairwise comparison matrices: (Quasi-)regular graphs with minimal diameter. *Omega*, 107, 102557.
- Tomashevskii, I. L. (2015). Eigenvector ranking method as a measuring tool: Formulas for errors. *European Journal of Operational Research*, 240(3), 774–780.
- Vaidya, O. S., & Kumar, S. (2006). Analytic hierarchy process: An overview of applications. *European journal of Operational Research*, 169(1), 1–29.
- Vargas, L. G. (1982). Reciprocal matrices with random coefficients. *Mathematical Modelling*, 3(1), 69–81.
- Vargas, L. G. (1990). An overview of the analytic hierarchy process and its applications. *European Journal of Operational Research*, 48(1), 2–8.
- Williams, C., & Crawford, G. (1980). Analysis of subjective judgment matrices. *Interim report R-2572-AF*. Santa Monica: Rand Corporation.